# **B.1** Hydrologic Region Profile: North Coast (NC)



## B.1.1 Summary

The North Coast Hydrologic Region is in the northwestern corner of California and includes redwood forests, inland mountain valleys, and the arid Modoc Plateau. Heavy rainfall in the coastal mountain ranges makes the North Coast


region the most water-abundant area of California, producing about 41 percent of the state's total natural runoff. As a result of the abundant rainfall, the average annual runoff for the rivers of this region is about 29 million acre-feet per year, which is the largest volume compared to all other hydrologic regions of California.

See Table 1 for additional information on the North Coast Hydrologic Region.

Size	19,476 square miles (2	12.3% of State)			
Population (2000)	644,000				
<b>Projected Population</b>	895,150		Percent Growth	39%	
(2030)					
Reservoir Storage	3,780 TAF				
Capacity					
Average Annual	50.6 inches				
Precipitation					
Land Use	Forest and rangeland	represent about	98 percent of this reg	ion's land area.	
	The agricultural trend	in the past deca	de has been one of la	nd consolidation	
	and the conversion of	prime agricultu	ral land to urban grow	th. This trend is a	
	result of low crop values, the lack of additional inexpensive surface water,				
	and the ability to use	only the most ec	conomically developab	le groundwater.	
Water Supply	Many of the smaller communities and rural areas in the North Coast region				
	are generally supplied by small local surface water and groundwater				
	systems. Larger water supply projects in this region include the USBR's				
	Klamath Project, the U.S. Army Corps of Engineers' Russian River Project				
	(Lake Mendocino and Lake Sonoma), and the Humboldt Bay Municipal Water				
	District's Ruth Reserve	oir, which serves	coastal communities	from Eureka to	
	McKinleyville.				
Water Use	In the year 2000, a "n	ormal" water ye	ar, agriculture accoun <sup>.</sup>	ts for about 83	
	percent of the region'	s water use, whi	le urban use is about :	17 percent.	
	Urban	126.9 TAF			
	Agricultural	640.7 TAF			
	Total	767.6 TAF			

Table 1 - North Coast Hydrologic Region Profile

## **B.1.2** North Coast 2010 Water Supply and Energy Profile:



#### Total Supply: 3,251 TAF

Total Energy Consumption: 159 GWh

# B.2 Hydrologic Region Profile: San Francisco Bay (SF)



## B.2.1 Summary

The San Francisco Bay Hydrologic Region, which occupies parts of nine counties, extends from southern Santa Clara County north to Tomales Bay in Marin County, and inland to the confluence of the Sacramento and San Joaquin rivers near Collinsville. The climate within the region varies significantly from


west to east. Coastal areas are typically cool and often foggy, and inland valleys are warmer with a Mediterranean-like climate.

See Table 2 for additional information on the San Francisco Bay Hydrologic Region.

		<u> </u>	-9		
Size	4,506 square feet (2.85 of State)				
Population (2000)	6,105,650				
Projected Population	7,857,360		Percent Growth	29%	
(2030)					
Reservoir Storage	746 TAF				
Capacity					
Average Annual	25.4 inches				
Precipitation					
Land Use	Portions of the region are highly urbanized and include the San Francisco,				
	Oakland, and San Jose	e metropolitan ai	reas. Agricultural acrea	ge occurs	
	mostly in the north ar	nd northeast in N	lapa, Marin, Sonoma, a	ind Solano	
	counties. Santa Clara	and Alameda co	unties also have signifi	cant agricultural	
	acreage at the edge o	lge of the urban development.			
Water Supply	In the early 1900s, loc	al water agencie	s developed significant	t imported	
	water supplies from tl	ne Mokelumne a	nd Tuolumne rivers to	meet the	
	anticipated demands.	At the same tim	e, local reservoirs and	watersheds	
	were being developed	l to capture surfa	ace supplies, to rechar	ge the	
	groundwater basins, a	and to act as terr	ninal reservoirs for the	larger projects.	
	Recycled water is also	used in a full sp	ectrum of applications	including	
	landscape irrigation, in	ndustrial cooling	, agricultural needs, an	d as supply to	
	wetlands. Currently, I	nearly 50-million	gallons per day of recy	cled water is	
	produced in the bay re	egion.			
Water Use	In the year 2000, a "n	ormal" water yea	ar, agriculture accounts	s for about 10%	
	percent of the region'	s water use, whi	le urban use is about 9	0% percent.	
	Urban	1,011.8 TAF			
	Agricultural	109.7 TAF			
	Total	1,121.5 TAF			

Table 2 – San Francisco Bay Hydrologic Region Profile

## B.2.2 San Francisco Bay 2010 Water Supply and Energy Profile:



#### Total Supply: 1,277 TAF

**Total Energy Consumption: 103 GWh** 

# **B.3 Hydrologic Region Profile: Central Coast (CC)**



### B.3.1 Summary

The Central Coast Hydrologic Region extends from southern San Mateo County in the north to Santa Barbara County in the south. The region includes all of Santa Cruz, Monterey, San Benito, San Luis Obispo and Santa Barbara counties and parts of San Mateo, Santa Clara, and Ventura counties. Many attributes

define the Central Coast region including: the topography, many microclimates, the variety of agricultural products, and the picturesque coastline, Valleys and communities that drive a thriving tourism economy.


See Table 3 for additional information on the Central Coast Hydrologic Region.

	Table 3 – Central Coas	t Hydrologic Reg	ion Profile			
Size	11,326 square miles (7.1% of the state)					
Population (2000)	1,459,205					
<b>Projected Population</b>	1,890,390		Percent Growth	30%		
(2030)						
Reservoir Storage	1,227 TAF					
Capacity						
Average Annual	18.7 inches					
Precipitation						
Land Use	The busy topography	of the Central C	oast Region and dista	nce from		
	California's major pop	oulation centers	have resulted in a lan	dscape that is		
	primarily pastoral and	l agricultural. Ag	riculture in the Centra	al Coast region		
	can be divided into tw	can be divided into two distinct categories. One is irrigated vegetable and				
	specialty crops grown on coastal terraces and valleys and in some inland					
	valleys; and the other category is range pasture and dry-farmed grain in the					
	inland valleys.					
Water Supply	Groundwater is the primary source of water in the region, accounting for					
	roughly 75 percent of the annual supply in 2000. Local and some imported					
	surface water supplie	•				
	region. A significant a	-		•		
	Pajaro, Salinas, and Ca	armel rivers, and	by the Arroyo Seco,	which flows into		
	the Salinas river.					
Water Use	In the year 2000, a "n		· · ·			
	percent of the region		ile urban use is about	23% percent.		
	Urban	221.0 TAF				
	Agricultural	746.2 TAF				
	Total 967.2 TAF					


#### Table 3 – Central Coast Hydrologic Region Profile

### B.3.2 Central Coast 2010 Water Supply and Energy Profile:

Total Supply: 1,219 TAF



#### Total Energy Consumption: 435 GWh



# B.4 Hydrologic Region Profile: South Coast (SC)



## B.4.1 Summary

The South Coast Hydrologic Region comprises the southwest portion of the state and is California's most urbanized and populous region. The topography includes a series of nearly flat coastal plains and valleys, many broad but gentle interior valleys, and several mountain ranges of low and moderate elevation.


The region has a mild, dry subtropical climate where summers are virtually rainless, except in the mountains where late summer thunderstorms sometimes occur.

See Table 4 for additional information on the South Coast Hydrologic Region.

Size	10.025 square miles (	, , ,				
	10,925 square miles (6.9% of State)					
Population (2000)	18,223,425					
Projected Population	23,827,075		Percent Growth	24%		
(2030)						
Reservoir Storage	3,059 TAF					
Capacity						
Average Annual	17.6 inches					
Precipitation						
Land Use	The expansion of new	single- and mul	ti-family homes, comm	ercial services,		
	businesses, and highv	vay systems into	the warmer sections of	f the region		
	continues onto lands	that were histor	ically pastoral, if not ag	ricultural.		
			agricultural uses still exi			
		· ·	entura County to the int			
			, m the coast to beyond I			
			•			
	San Bernardino. Irrigated agriculture now occupies only one-seventh as much land as urban uses.					
Water Supply	The region has developed a diverse mix of both local and imported water					
	supply sources. Local water resources development over the last 15 years					
			•	•		
		has included water recycling, groundwater storage and conjunctive use, conservation, brackish water desalination, water transfer and storage, and				
			plement imported wate	•		
		•	te Water Project (SWP),			
		-	geles Aqueduct (LAA). T			
		-	ging supplies and resou			
	dry years.	,	00			
Water Use		ormal" water ve	ar, agriculture accounts	s for about 17%		
			ile urban use is about 8			
	Urban	3,860.4 TAF				
	Agricultural	795.9 TAF				
	Total 4656.3 TAF					
	10tal 4050.3 TAF					

#### Table 4 – South Coast Hydrologic Region Profile

## B.4.2 South Coast 2010 Water Supply and Energy Profile:



#### Total Supply: 5,408 TAF



Note: the negative area is due to energy production in the region

# **B.5** Hydrologic Region Profile: Sacramento River (SR)



## B.5.1 Summary

The Sacramento River Hydrologic Region includes the entire drainage area of the state's largest river and its tributaries, extending from the Oregon border downstream to the Sacramento – San Joaquin Delta. The Sacramento Valley with its alluvial soils, abundant water and moderate climate, is one of the richest


agricultural regions on earth. These same physical attributes also make it an incredibly productive ecosystem that supports more than 250 species of fish and wildlife.

See Table 5 below for additional information on the Sacramento River Hydrologic Region.

	Table 5 - Sacramento Ki	, ,	egion i rome		
Size	27,246 square miles (	17.2% of State)			
Population (2000)	2,593,110				
Projected Population	4,569,490		Percent Growth	76%	
(2030)					
Reservoir Storage	16,146 TAF				
Capacity					
Average Annual	36.7				
Precipitation					
Land Use	Agriculture is the regi	on's largest indu	stry, contributing a wic	le	
	variety of crops includ	ding rice, grain, t	omatoes, field crops, fr	uits	
	and nuts. A substantia	al number of acr	es of rangeland in this r	egion are also	
	used for livestock ma	nagement. Durin	ng the past 130 years, m	nore than 95	
	percent of the valley's	s historic ripariar	n forests have been cor	verted to other	
	land uses.				
Water Supply	Because of the weather patterns that produce a high level of precipitation in				
	the region, major wat	er supplies from	the region are provide	d through the	
	development of reser	voirs and from d	lirect groundwater pur	nping, which	
	historically has rechar	ged through the	winter months. Althou	igh a few of the	
	larger cities in the reg	ion, such as Saci	ramento, divert most of	f their water	
	from the larger rivers	, the principal so	urce of water for most	of the urban	
	and rural communitie	s throughout thi	s region is groundwate	r.	
Water Use	In the year 2000, a "n	ormal" water ye	ar, agriculture accounts	s for about 91%	
	percent of the region	's water use, whi	ile urban use is about 9	% percent.	
	Urban	756.2 TAF			
	Agricultural	7,343.3 TAF			
	Total	8,099.5 TAF			

Table 5 – Sacramento River Hydrologic Region Profile

## B.5.2 Sacramento River 2010 Water Supply and Energy Profile:



### Total Supply: 21,545 TAF

#### Total Energy Consumption: 1,068 GWh

# B.6 Hydrologic Region Profile: San Joaquin River (SR)



### B.6.1 Summary

The San Joaquin River hydrologic region is in the heart of California and includes the northern portion of the San Joaquin Valley. It is bordered on the east by the Sierra Nevada and on the west by the coastal mountains of the Diablo Range. Because the San Joaquin Valley is isolated by mountains from


the marine effects of coastal California, the maximum average daily temperature in the valley reaches a high of 101 degrees during late July, although the northern part of this hydrologic region does benefit from Delta breezes during the hot summer periods.

See Table 6 below for additional information on the San Joaquin River Hydrologic Region.

	lable 6 – San Joaquin Ri		egion Profile		
Size	15,214 square miles (9.6% of State)				
Population (2000)	1,751,010				
Projected Population	3,385,885		Percent Growth	93%	
(2030)					
Reservoir Storage	11,477 TAF				
Capacity					
Average Annual	26.3 inches				
Precipitation					
Land Use	The valley portion of t	he San Joaquin I	River region consists p	rimarily of highly	
	productive farmland a	and the rapidly g	rowing urban areas of	Stockton, Tracy,	
	Modesto, Manteca, a	nd Merced. Agrid	culture is the major ec	onomic and land	
	use activity in the San Joaquin River region. The Valley has roughly 2 million				
	acres of irrigated cropland and an annual agricultural output valued at more				
	than \$ 4.9 billion.				
Water Supply	The primary sources of surface water in the San Joaquin River region are the				
	rivers that drain the w	•			
	San Joaquin River and	-			
	Stanislaus, Calaveras,				
	water in the upper Sa	•			
	and is then conveyed	north through th	ne Madera Canal and s	south through	
	the Friant-Kern Canal.				
Water Use	In the year 2000, a "n	-	-		
	percent of the region'	s water use, whi	le urban use is about	6% percent.	
	Urban	374.1 TAF			
	Agricultural	5655.9 TAF			
	Total	6,030.0 TAF			

Table 6 – San Joaquin River Hydrologic Region Profile

## 1.1.1 San Joaquin River 2010 Water Supply and Energy Profile:



#### Total Supply: 10,448 TAF

#### Total Energy Consumption: 2,682 GWh

# 1.2 Hydrologic Region Profile: Tulare Lake (TL)



### 1.2.1 Summary

The Tulare Lake Hydrologic Region is in the southern end of the San Joaquin Valley. This region includes all of Tulare and Kings Counties and large portions of Fresno and Kern counties. The valley is broad and flat, and is surrounded by the Diablo and Coast Ranges to the west, the Sierra Nevada to the east, and the


Tehachapi Mountains to the south. The valley portion of the region is hot and dry in summer with long, sunny days and cooler nights. Winters are wet and often blanketed with dense fog.

See Table 7 below for additional information on the Tulare Lake Hydrologic Region.

	Table 7 – Tulare Lake Hydrologic Region Profile				
Size	17,033 square miles (2	10.7% of State)			
Population (2000)	1,884,675				
Projected Population	3,121,625		Percent Growth	66%	
(2030)					
Reservoir Storage	2,046 TAF				
Capacity					
Average Annual	15.2 inches				
Precipitation					
Land Use	The State and federal	government age	encies own about 30 pe	rcent of the	
	land in the region, inc	luding about 1.7	million acres of nation	al forest, 0.8	
	million acres of natior	nal parks and rec	reation areas, and 1 mi	illion acres of	
	land managed by the	U.S. Bureau of La	and Management. Priva	ately owned	
	land totals about 7.4 million acres. Irrigated agriculture accounts for more				
	than 3 million acres of the private land, while urban areas take up over				
	350,000 acres. Other agricultural lands and areas with native vegetation				
	represent an addition	al 1.4 million acr	es in the region.		
Water Supply	The region receives most of its surface water runoff from four main rivers				
	that flow out of the Sierra Nevada, which are the Kings, Kaweah, Tule, and				
	Kern rivers. Major wat	ter conveyance f	acilities in the region in	nclude the	
	California Aqueduct, t	he Friant-Kern C	anal, and the Cross Val	ley Canal.	
	Water diversions from	n the San Joaquii	n River at Friant Dam ar	re also a	
	significant supply sour	rce for all uses in	the Tulare Lake region	l.	
Water Use	In the year 2000, a "n	ormal" water ye	ar, agriculture accounts	s for about 97%	
	percent of the region'	s water use, whi	le urban use is about 3	% percent.	
	Urban	236.1 TAF			
	Agricultural	8,104.8 TAF			
	Total	8340.9 TAF			


#### Table 7 – Tulare Lake Hydrologic Region Profile

## B.6.2 Tulare Lake 2010 Water Supply and Energy Profile:



### Total Supply: 11,258 TAF To

### Total Energy Consumption: 6,736 GWh



# B.7 Hydrologic Region Profile: North Lahontan (NL)



## B.7.1 Summary

The North Lahontan Hydrologic Region forms part of the western edge of the Great Basin, a large landlocked area that includes most of Nevada and northern Utah. The crest of the Sierra Nevada forms much of the western boundary of this region. All surface water in the region drains eastward toward Nevada.


The regions climate is characterized by dry summers, with the occasional thunderstorm and snowy winters. The snowpack from the eastern slopes of the Sierra Nevada melts in the late spring and summer to become the primary source of surface water supplies for much of northern Nevada.

See Table 8 below for additional information on the North Lahontan Hydrologic Region.

	Table 8 – North Lahonta	, ,			
Size	6,122 square miles (3.9% of the state)				
Population (2000)	99,035				
Projected Population	130,800		Percent Growth	32%	
(2030)					
Reservoir Storage	1,181 TAF				
Capacity					
Average Annual	23.1 inches				
Precipitation					
Land Use	Much of the region is	either national f	orest or lands under th	ne jurisdiction of	
	the Bureau of Land M	anagement. Cat	tle-ranching is the prin	cipal agricultural	
	activity with pasture a	and alfalfa being	the dominant irrigated	d crops.	
	Although growing sea	sons vary consid	erably each year, the r	nountain valleys	
	where most crops are	grown are usua	lly frost free from late	May to mid-	
	September or about 1	.20 days.			
Water Supply	The largest rivers in th	ne region and the	eir average regulated r	unoff at the	
	Nevada state line are	the Truckee Rive	er with 540,000 acre-fe	et; the Carson	
	River, 335,000 acre-fe	et; and the Wall	ker River, 300,000 acre	-feet. The Susan	
	River is the only majo	r river in the nor	thern half of the region	n and its annual	
	discharge at Susanville	e averages 60,00	0 acre-feet. The Truck	kee, Carson, and	
	Walker rivers are large	ely governed by	existing federal court v	water right	
	decrees. On the Califo	ornia side, most l	ocally developed wate	r supplies are	
	from groundwater or	small surface wa	ater diversions, with st	orage provided	
	by outlet dams constr	ucted on natura	l lakes.		
Water Use	In the year 2000, a "n	ormal" water ye	ar, agriculture account	s for about 94%	
	percent of the region'	s water use, whi	le urban use is about 6	5% percent.	
	Urban	24.8 TAF			
	Agricultural	400.1 TAF			
	Total 424.9 TAF				

 Table 8 – North Lahontan Hydrologic Region Profile

## B.7.2 North Lahontan 2010 Water Supply and Energy Profile:



#### Total Supply: 691 TAF

#### **Total Energy Consumption: 49 GWh**

# 1.3 Hydrologic Region Profile: South Lahontan (SL)



### 1.3.1 Summary

The South Lahontan Hydrologic Region contains the Eastern Sierra and the Mojave Desert and includes both the highest point (Mount Whitney) and lowest point (Death Valley) in the lower 48 states. The northern half of the region includes Mono Lake, Owens Valley, Panamint Valley, Death Valley,

and the Amargosa River Valley. The climate of the South Lahontan region is generally arid.


See Table 9Error! Reference source not found. below for additional information on the South Lahontan Hydrologic Region.

Size		, ,	J	26,732 square miles (16.9% of State)			
Population (2000)	721,490						
Projected Population	1,266,375		Percent Growth	76%			
(2030)	1,200,575		r creent Growth	7070			
	459 TAF						
Reservoir Storage	459 TAF						
Capacity	7011						
Average Annual	7.8 inches						
Precipitation							
Land Use	The region supports a variety of urban and agricultural uses, including a						
	moderate amount of	agricultural acre	age and several growi	ng cities. Much			
	of the land in the region remains undeveloped and is under protected or						
	managed status for recreational, scenic, environmental, or military purposes.						
Water Supply	The Los Angeles Aque	The Los Angeles Aqueduct is the region's major water development feature.					
	In 1913, the initial 223	In 1913, the initial 223-mile-long aqueduct was completed by the Los					
	Angeles Department of Water and Power (LADWP) and began transporting						
	water from Owens Valley to the city of Los Angeles. Since then the aqueduct						
	has been extended 115 miles and a second 137-mile pipeline was						
	completed. There are						
	system with a combin	-	-				
Water Use	In the year 2000, a "n		-				
Water Ose	percent of the region'						
	· · ·		ile ul ball use is about	5770 percent.			
	Urban	178.8 TAF					
	Agricultural	310.4 TAF					
	Total	489.2 TAF					


Table 9 – South Lahontan Hydrologic Region Profile

B.7.3 South Lahontan 2010 Water Supply and Energy Profile:

### Total Supply: 730 TAF



### Total Energy Consumption: 657 GWh



# B.8 Hydrologic Region Profile: Colorado River (CR)



### B.8.1 Summary

The Colorado River Hydrologic Region is in the southeastern corner of California. The region includes all of Imperial County, about the eastern onefourth of San Diego County, the eastern two-thirds of Riverside County, and the southeastern one-third of San Bernardino County. It has a variety of arid desert terrain that includes many bowl-shaped valleys, broad alluvial fans,


sandy washes, and hills and mountains. Nearly the entire Colorado River region has a subtropical desert climate with hot summers and generally mild winters.

See Table 10 below for additional information on the Colorado River Hydrologic Region.

Size	Table 10 – Colorado River Hydrologic Region Profile				
	19,962 square miles (12.6% of State)				
Population (2000)	606,535				
Projected Population	1,166,550		Percent Growth	92%	
(2030)					
Reservoir Storage	620 TAF				
Capacity					
Average Annual	5.7 inches				
Precipitation					
Land Use	The region is a land of	f unequalled agr	icultural bounty with a	a growing urban	
	sector, and large expa	anses of open, w	ild terrain. Famous pa	rks in the region	
	include Joshua Tree N	lational Park, the	e Mojave National Sce	nic Preserve,	
	Anza-Borrego Desert	State Park, and t	the Salton Sea and the	Picacho state	
	recreation areas. The	re are also sever	al areas set aside for p	preservation or	
	other land manageme	ent purposes, ind	cluding national recrea	ation and	
	wilderness areas, wild	llife refuges, Ind	ian tribal reservations	and U.S. Navy	
	facilities.				
Water Supply	About 85 percent of the region's urban and agricultural water supply comes				
	from surface water de	eliveries from th	e Colorado River. Wat	er from the river	
	is delivered into the r	egion through th	ne All-American and Co	oachella canals,	
			er Aqueduct by means		
			The remaining water i	-	
	-		The Colorado River is	•	
		•	pportioned among the		
			mplex body of statute		
			the "Law of the River"		
Water Use			ar, agriculture accoun		
			ile urban use is about		
	Urban	420.6 TAF			
	Agricultural	3,831.6 TAF			
Total 4,252.2 TAF					
	10ldi 4,252.2 IAF				

Table 10 – Colorado River Hydrologic Region Profile

## B.8.2 Colorado River 2010 Water Supply and Energy Profile:



### Total Supply: 4,697 TAF

Total Energy Consumption: 2,025 GWh